$[ BoldFont = LinLibertine_RB.otf, ItalicFont = LinLibertine_RI.otf, BoldItalicFont = LinLibertine_RBI.otf, Path = /opt/indico/.venv/lib/python2.7/site-packages/indico_fonts/] [BoldFont = LinBiolinum_RB.otf, ItalicFont = LinBiolinum_RI.otf, Path = /opt/indico/.venv/lib/python2.7/site-packages/indico_fonts/] [BoldFont = LinBiolinum_RB.otf, ItalicFont = LinBiolinum_RI.otf, Path = /opt/indico/.venv/lib/python2.7/site-packages/indico_fonts/] [BoldFont = LinBiolinum_RB.otf, ItalicFont = LinBiolinum_RI.otf, Path = /opt/indico/.venv/lib/python2.7/site-packages/indico_fonts/] [BoldFont = LinBiolinum_RB.otf, ItalicFont = LinBiolinum_RI.otf, Path = /opt/indico/.venv/lib/python2.7/site-packages/indico_fonts/] [BoldFont = LinBiolinum_RB.otf, ItalicFont = LinBiolinum_RI.otf, Path = /opt/indico/.venv/lib/python2.7/site-packages/indico_fonts/] [BoldFont = LinBiolinum_RI.otf, Path = /opt/indico/.venv/lib/python2.7/site-packages/indico_fonts/] [Bol$ 

HTPD 2018



Contribution ID : 215

Type : not specified

## 4.34 DIII-D Faraday-Effect Polarimeter for Fast Magnetic Dynamics Measurement

Monday, 16 April 2018 20:31 (120)

A Radial-Interferometer-Polarimeter (RIP) diagnostic has been developed to explore fast magnetic dynamics in high-performance DIII-D plasmas by measuring radial magnetic field perturbations using three chords positioned near the magnetic axis. Newly developed solid-state sources operated at 650 GHz are used and provide phase noise down to 0.01 degree/sqrt(kHz) and tunable bandwidth up to 10 MHz. Various systematic errors, which can contaminate the Faraday-effect polarimetric measurement, have been investigated in detail. The impact of the perpendicular magnetic field (Cotton-Mouton Effect) is evaluated and found negligible. Distortion of circular polarization due to non-ideal optical components is calibrated by using a rotating quarter wave-plate technique. Optical feedback, due to multiple reflections induced by the double-pass configuration, is reduced by increasing the damping of stray light. Error due to non-collinearity is minimized to less than 0.5 degree for electron density up to 1x1020m-3 by optimizing the alignment. Measurement of coherent and broadband high-frequency magnetic fluctuation for DIII-D H-mode plasmas is presented. This material is based upon work supported by the Department of Energy under Award Numbers DE-FG03-01ER54615 and DE-FC02-04ER54698.

Primary author(s): CHEN, Jie (Physics & Astronomy, UCLA )

Co-author(s) : DING, Weixing (Physics & Astronomy, UCLA ); BROWER, David (Physics & Astronomy, UCLA ); FINKENTHAL, Daniel (Palomar Scientific Instruments); BOIVIN , Rejean (General Atomics )

Presenter(s) : CHEN, Jie (Physics & Astronomy, UCLA ); DING, Weixing (Physics & Astronomy, UCLA ); BROWER, David (Physics & Astronomy, UCLA ); FINKENTHAL, Daniel (Palomar Scientific Instruments); BOIVIN , Rejean (General Atomics )

Session Classification : Session #4, Monday Night Poster Session