

HTPD 2018



Contribution ID : 247

Type : not specified

## 8.3 OMEGA Supersonic Gas-Jet Plasma Characterization with Thomson Scattering

Tuesday, 17 April 2018 16:00 (120)

A supersonic gas-jet target platform has been activated on the OMEGA laser. The plasma formed using a gas-jet target and ~3 kJ of UV energy from the OMEGA Laser System was characterized using  $2\omega$  Thomson scattering. Thomson scattering provided accurate time-resolved measurements of plasma conditions including electron density, plasma temperature, and ionization state. Plasma conditions include electron temperatures in the 0.5-keV to 1-keV range and electron densities between  $1 \times 10^{19} \text{cm}^{-3}$  and  $9 \times 10^{19} \text{cm}^{-3}$  in a nitrogen plasma. The measurements made using Thomson scattering are then compared with the results of the radiation-hydrodynamics code HYDRA. These initial measurements demonstrate the capabilities of the OMEGA gas-jet as a platform for future laser-plasma interaction science. This material is based upon work supported by the Department of Energy National Nuclear Security Administration under Award Number DE-NA0001944.

Primary author(s) : HANSEN, Aaron (Laboratory for Laser Energetics)

Co-author(s) : HABERBERGER, Daniel (Laboratory for Laser Energetics); FROULA, Dustin (Laboratory for Laser Energetics)

Presenter(s) : HANSEN, Aaron (Laboratory for Laser Energetics); HABERBERGER, Daniel (Laboratory for Laser Energetics); FROULA, Dustin (Laboratory for Laser Energetics)

Session Classification : Session #8, Tuesday Afternoon Poster Session