

HTPD 2018



Contribution ID : 323

Type : not specified

## 8.24 A new method to reconstruct the Bp profile in the Laser-driven Ion-beam Trace Probe (LITP) diagnostics

Tuesday, 17 April 2018 16:01 (120)

Xiaoyi Yang, Tianchao Xu, Yihang Chen, Tianbo Wang, Chijie Xiao, Min Xu, Yi Yu, Xiaogang Wang  
The Laser-driven Ion-beam Trace Probe (LITP) is a new Bp diagnostic method, firstly proposed in 2014. The basic principle of the LITP method is as follow. Ions generated by laser-driven accelerator are injected into the tokamak, passing through the plasma and finally reached the detector on the vacuum wall. The Bp profile could be reconstructed based on the traces of ions. The ion beam has large energy and pitch angle spread so that the traces of ions with different energies cover a 2D area in the tokamak, and a tomography method is needed to reconstruct 2D Bp profiles. Here a new method based on the solutions of differential equation is proposed. Simulation results and error analysis show that the new method makes the LITP more adjustable and robust.

[1] Yang et al. Rev. Sci. Instrum. 85(11), 11E429 (2014).

[2] Yang et al. Rev. Sci. Instrum. 87(11), 11D610 (2016).

Primary author(s) : YANG, Xiaoyi (Peking University)

Presenter(s) : YANG, Xiaoyi (Peking University)

Session Classification : Session #8, Tuesday Afternoon Poster Session