

HTPD 2018



Contribution ID : 386

Type : not specified

6.9 Synthetic Diagnostic for Assessing Spatial Averaging of Charge Exchange Recombination Spectroscopy Measurements

Tuesday, 17 April 2018 10:30 (120)

A synthetic charge exchange recombination spectroscopy diagnostic based on the FIDAsim modeling suite has been created for the DIII-D tokamak. This synthetic diagnostic assumes the ions have Maxwellian distribution functions on each flux surface and generates synthetic emission from charge exchange events between the beam neutrals and a fully ionized impurity. This work was motivated by the observation of non-Gaussian spectra that may be caused by spatial averaging, atomic physics, or non-Maxwellian distribution functions. Measurements of non-Gaussian spectra commonly observed in the H-mode pedestal and in plasmas with very steep core gradients are compared to the synthetic diagnostic. Spatial averaging alone cannot account for the observations, indicating other cause(s) such as non-Maxwellian distribution functions. The synthetic diagnostic has also been used to resolve a long-standing issue: it is shown that vertical view chords near the magnetic axis often measure lower temperatures than the tangential view chords because of a difference in spatial averaging due to the DIII-D neutral beams being twice as tall as they are wide. Work supported by US DOE under DE-FC02-04ER54698, DE-AC02-09CH11466, and the Science Undergraduate Laboratory Internship (SULI) program.

Primary author(s) : CHRYSTAL, Colin (General Atomics)

Co-author(s) : SULYMAN, Alex (University of California Davis); HASKEY, Shaun (Princeton Plasma Physics Laboratory); BURRELL, Keith (General Atomics)

Presenter(s) : CHRYSTAL, Colin (General Atomics); SULYMAN, Alex (University of California Davis); HASKEY, Shaun (Princeton Plasma Physics Laboratory); BURRELL, Keith (General Atomics); GRIERSON, Brian (Princeton Plasma Physics Laboratory)

Session Classification : Session #6, Tuesday Morning Poster Session