$[ BoldFont = LinLibertine_RB.otf, ItalicFont = LinLibertine_RI.otf, BoldItalicFont = LinLibertine_RBI.otf, Path = /opt/indico/.venv/lib/python2.7/site-packages/indico_fonts/] [BoldFont = LinBiolinum_RB.otf, ItalicFont = LinBiolinum_RI.otf, Path = /opt/indico/.venv/lib/python2.7/site-packages/indico_fonts/] [BoldFont = LinBiolinum_RB.otf, ItalicFont = LinBiolinum_RI.otf, Path = /opt/indico/.venv/lib/python2.7/site-packages/indico_fonts/] [BoldFont = LinBiolinum_RB.otf, ItalicFont = LinBiolinum_RI.otf, Path = /opt/indico/.venv/lib/python2.7/site-packages/indico_fonts/] [BoldFont = LinBiolinum_RB.otf, ItalicFont = LinBiolinum_RI.otf, Path = /opt/indico/.venv/lib/python2.7/site-packages/indico_fonts/] [BoldFont = LinBiolinum_RB.otf, ItalicFont = LinBiolinum_RI.otf, Path = /opt/indico/.venv/lib/python2.7/site-packages/indico_fonts/] [BoldFont = LinBiolinum_RI.otf, Path = /opt/indico/.venv/lib/python2.7/site-packages/indico_fonts/] [Bol$ 

**HTPD 2018** 



Contribution ID : 415

Type : not specified

## 6.36 Developing a bright high-energy continuum backlighter for EXAFS on NIF

Tuesday, 17 April 2018 10:31 (120)

Extended X-ray Absorption Fine Structure (EXAFS) spectroscopy is a powerful tool for in situ characterization of matter in the high energy density regime. An EXAFS platform is currently being developed on the National Ignition Facility (NIF). Development of a suitable X-ray backlighter involves minimizing the temporal duration and source size while maximizing spectral smoothness and brightness. One approach involves imploding a spherical shell, which generates a large of amount of heat and an X-ray flash near stagnation. Radiation hydrodynamics modeling improvements for EXAFS are possible by filling the shell with a moderate-Z gas. Here we present measurements of X-ray source size, spectral-temporal emission, and integrated spectrum produced by imploded Ar-filled CH shells. Compared to an unfilled shell, we find that 1 atm Ar fill significantly increases the X-ray yield but also increases the source size, whereas 4 atm Ar fill produces a similar yield but reduced the source size. This work performed under the auspices of the Lawrence Livermore National Security, LLC, (LLNS) under Contract No. DE-AC52-07NA27344.

Primary author(s): KRYGIER, Andrew (Lawrence Livermore National Laboratory)

Co-author(s): COPPARI, Federica (Lawrence Livermore National Laboratory); KEMP, Gregory Elijah (Lawrence Livermore National Laboratory); THORN, Daniel (Lawrence Livermore National Laboratory); EGGERT, Jon (Lawrence Livermore National Laboratory); PARK, Hye-Sook (Lawrence Livermore National Laboratory); PING, Yuan (Lawrence Livermore National Laboratory); SCHNEIDER, Marilyn (Lawrence Livermore National Laboratory)

Presenter(s): KRYGIER, Andrew (Lawrence Livermore National Laboratory); COPPARI, Federica (Lawrence Livermore National Laboratory); KEMP, Gregory Elijah (Lawrence Livermore National Laboratory); THORN, Daniel (Lawrence Livermore National Laboratory); EGGERT, Jon (Lawrence Livermore National Laboratory); PARK, Hye-Sook (Lawrence Livermore National Laboratory); PING, Yuan (Lawrence Livermore National Laboratory); REM-INGTON, Bruce (Lawrence Livermore National Laboratory); SCHNEIDER, Marilyn (Lawrence Livermore National Laboratory)

Session Classification : Session #6, Tuesday Morning Poster Session