$[BoldFont = LinLibertine_RB.otf, ItalicFont = LinLibertine_RI.otf, BoldItalicFont = LinLibertine_RBI.otf, Path = /opt/indico/.venv/lib/python2.7/site-packages/indico_fonts/] [BoldFont = LinBiolinum_RB.otf, ItalicFont = LinBiolinum_RI.otf, Path = /opt/indico/.venv/lib/python2.7/site-packages/indico_fonts/] [BoldFont = LinBiolinum_RB.otf, ItalicFont = LinBiolinum_RI.otf, Path = /opt/indico/.venv/lib/python2.7/site-packages/indico_fonts/] [BoldFont = LinBiolinum_RB.otf, ItalicFont = LinBiolinum_RI.otf, Path = /opt/indico/.venv/lib/python2.7/site-packages/indico_fonts/] [BoldFont = LinBiolinum_RB.otf, ItalicFont = LinBiolinum_RI.otf, Path = /opt/indico/.venv/lib/python2.7/site-packages/indico_fonts/] [BoldFont = LinBiolinum_RB.otf, ItalicFont = LinBiolinum_RI.otf, Path = /opt/indico/.venv/lib/python2.7/site-packages/indico_fonts/] [BoldFont = LinBiolinum_RI.otf, Path = /opt/indico/.venv/lib/python2.7/site-packages/indico_fonts/] [Bol$

HTPD 2018



Contribution ID : 417

Type : not specified

6.38 Characterization of Selenium He x-ray source on the National Ignition Facility

Tuesday, 17 April 2018 10:31 (120)

There are many high-energy-density experiments that require efficient atomic line emission x-ray sources for diagnostic applications such as imaging (e.g. backlit radiography) and material testing (e.g. diffraction measurements). To date, most well-characterized laser-generated line sources efficient enough for these purposes have photon energies <10.2 keV. They are typically created by irradiating a thin foil using a 351nm, long pulse laser (\geq 1 ns) in the range of 1015 W/cm2. The dominant line emission, He α , from these sources is the result of 2p -> 1s transitions from He-like ions. For the new Crystal Backlighter Imager at the National Ignition Facility (NIF), we developed a Selenium He α source at 11.652 keV. The Se He-like line emission was investigated in terms of absolute spectra and laser conversion efficiency into the lines as a function of viewing angle relative to the foil normal. Time-integrated and time-resolved data from multiple NIF shots will be presented. This work was performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344 and by General Atomics under Contract DE-NA0001808.

Primary author(s) : KRAULAND, Christine (General Atomics)

Co-author(s) : BARRIOS, Maria A. (Lawrence Livermore National Laboratory); THORN, Daniel B. (Lawrence Livermore National Laboratory); SCHNEIDER, Marilyn B. (Lawrence Livermore National Laboratory); HALL, Gareth N. (Lawrence Livermore National Laboratory); LANDEN, Otto L. (Lawrence Livermore National Laboratory)

Presenter(s) : KRAULAND, Christine (General Atomics); BARRIOS, Maria A. (Lawrence Livermore National Laboratory); THORN, Daniel B. (Lawrence Livermore National Laboratory); SCHNEIDER, Marilyn B. (Lawrence Livermore National Laboratory); HALL, Gareth N. (Lawrence Livermore National Laboratory); LANDEN, Otto L. (Lawrence Livermore National Laboratory)

Session Classification : Session #6, Tuesday Morning Poster Session