


Stability Control for Steady-State High-Beta in JT-60U




A. Isayama for the JT-60U team



Contents

JT-60U

- Introduction
 - Modification of JT-60U for long duration (65s) discharges
- Long-duration high- β_N plasma ($\beta_N=2.5$ for $\sim 10\tau_R$, $\beta_N=1.9$ for $\sim 15\tau_R$)
- High- β_N (~ 3) by NTM suppression
- Measurement of NTM structure
- Summary





Extension of discharge duration to 65s

JT-60U



Toroidal field : **3.3T for 30s, 2.7T for 65s**

Tangential P-NB (4 units): **30s injection** (<- formerly 10s)

N-NB: **~2MW for 30 s**




$\beta_N \sim 2.5$, $H_{89PL} \sim 2$ have been sustained for 15.5s ($\sim 9.5\tau_R$)



- $\beta_N \sim 2.5$ for 15.5s: $9.5\tau_R$, $\sim 78\tau_E$
 - $H_{89PL} \sim 2 \rightarrow \beta_N H_{89PL} / q_{95}^2 \sim 0.4$ ~ITER baseline scenario
 - ITB and ETB for $>10\text{s}$
 - Confinement degradation at $t \sim 17\text{s}$:
density rise & ETB degradation
- $\tau_R = \mu_0 \langle \sigma_{NC} \rangle a^2 / 12$
(D.R. Mikkelsen Phys. Fluids B 1 (1989) 333.)


Operational regime before 2003

JT-60U



New operational regime: $\beta_N=2-2.5$ for $>10\tau_R$

JT-60U



- **Region of sustained beta has been significantly extended.**


$\beta_N \sim 2.5$ (~advanced operation in ITER) for 16.5s, $\sim 10\tau_R$

$\beta_N \sim 1.9$ (~inductive operation in ITER) for 24s, $\sim 15\tau_R$

- Low collisionality & Larmor radius regime: $v^*/v^*_{ITER} \sim 3$, $\rho^*/\rho^*_{ITER} \sim 3$
- NTM in longtime scale: **not observed** so far (at $\beta_N \sim 2.5$)

New operational regime: $\beta_N=2-2.5$ for $>10\tau_R$

JT-60U



- **Region of sustained beta has been significantly extended.**

$\beta_N \sim 2.5$ (~advanced operation in ITER) for 16.5s, $\sim 10\tau_R$

$\beta_N \sim 1.9$ (~inductive operation in ITER) for 24s, $\sim 15\tau_R$

- Low collisionality & Larmor radius regime: $v^*/v^*_{ITER} \sim 3$, $\rho^*/\rho^*_{ITER} \sim 3$
- NTM in longtime scale: **not observed** so far (at $\beta_N \sim 2.5$)

Issue for reaching high beta region = NTM suppression

JT-60U

JT-60U : Two scenarios for NTM *suppression*

NTM avoidance : Optimization of $p(r)$ & $q(r)$

- Hybrid Scenario regime ($q_{95} \sim 4.5$)
 $q=1.5$ at the center, $q=2$ at $\rho > \sim 0.7$
- Low-q regime ($q_{95} \sim 2.2$)
 $q=1.5$ and 2 at $\rho > \sim 0.7$

This talk




NTM stabilization : Active suppression with EC wave



- Early ECCD
ECCD **before** the NTM onset
(or **just after**)
- Late ECCD
ECCD **after** the NTM saturation

Capability of various heating profile & plasma shape control in JT-60U



Highlights in 2004


High β regime ($\beta_N \sim 3$)
2nd harmonic X-mode EC wave
New attempt in JT-60U

Stationary high-beta with $\beta_N \sim 3$ for $\sim 4\tau_R$ at $q_{95} \sim 2.2$

JT-60U




6.2s ($4.1\tau_R$)

1MA/1.7T, $\delta \sim 0.4$





- $\beta_N=3$, $\beta_t=2.4\%$
- $\beta_N H_{89PL} / q_{95}^2 \sim 0.75$

- Low-q in NB phase
- $I_i \sim 0.8$ (4.3s) \rightarrow 0.7 (4.9s)
- $\beta_N / I_i \sim 4$





Early injection with fundamental O-mode EC wave

JT-60U




\dot{B}/f is more efficiently suppressed by the early injection



Precise adjustment of EC injection angle is required also in early injection.


NTM stabilization with early ECCD in $\beta_N \sim 3$ regime

JT-60U




- Complete stabilization at $\beta_N \sim 3$ ($\beta_p \sim 1.8$)
with smaller NB power than for late ECCD
- Confinement improvement by NTM stabilization
- Requirement for complete stabilization:
 - JECCD~JBS at $q=3/2$
 - High accuracy of ECCD location

Same as 1st O-mode




Successful NTM stabilization in $\beta_N \sim 3$ regime


JT-60U



☺ Successful
Recovery of β & H-factor




☹ Unsuccessful
Misaligned ECSD



Detailed measurement of NTM structure

NTM: Magnetic island formation




M-shaped amplitude profile

Recent observations in JT-60U


- Change in $J(r)$ during NTM growth/stabilization
- NTM structure in \tilde{T}_e during stabilization

MSE
ECE



Localized change in current density has been observed during NTM growth/suppression


JT-60U



Decrease in j due to NTM growth

Increase in j due to NTM stabilization

Decrease in j due to NTM reappearance




- If configuration & $\beta \sim \text{const}$, change in MSE signal = change in $j(r)$ → direct measurement of δj

Decrease/increase in bootstrap current due to NTM growth/suppression




\tilde{T}_e profile suggests temperature increase INSIDE island

JT-60U




Hypothesis

Temperature increase inside island by EC



Deformed M-shaped profile



The hypothesis well explains

- Decrease at inner-half region
- Increase at outer-half region
- Fast response time ($\ll \tau_R$)
- No asymmetry for misaligned ECCD

Details are under investigation

Reduced heat transport inside the island in LHD

Estimation of χ_{\perp} from cold pulse propagation

$$\chi_{\perp}^{\text{in}}/\chi_{\perp}^{\text{out}} \sim 1/10$$






FIG. 3. Comparison of temperature perturbation between experiment and simulation. The time evolutions of the measured (solid line) and the simulated (broken line) temperature perturbation at different radii in the same discharges as (a) Fig. 1 and (b) Fig. 2, respectively.


Similar asymmetry was observed during EC mirror scan.

JT-60U



EC before the NTM onset suggests more efficient deposition inside island.

JT-60U



- Hypothesis leads to "One peak in \tilde{T}_e = deposition inside island"
 - 'ECH/ECCD before NTM onset is more efficient'
 - Further investigation required

NTM onset physics is the most challenging

JT-60U



- 'NTM: Seed island formation by sawtooth and fishbone → growth'
- However, NTM can appear without clear MHD event

Spontaneous NTM

Example in JT-60U

- Reappearance of NTM at t=10.7s
- Constant NB power
- $\beta_N \sim 1.5$ (7.5s) → 1.67 (10.7s): small increase
- No sawtooth, no fishbones

Experimental & theoretical works required



Summary

JT-60U

Long-duration high- β_N discharge

- $\beta_N=2.5$ for $\sim 10\tau_R$, $\sim 80\tau_E$; $\beta_N=1.9$ for $\sim 15\tau_R$, $>100\tau_E$
- No NTM in long timescale at $\beta_N < \sim 2.5$ **Low v^* & ρ^* regime**

NTM suppression at $\beta_N \sim 3$ regime

- NTM avoidance at $q_{95} \sim 2.3$: **importance of $p(r)$ & $q(r)$**
- NTM stabilization by early ECCD:
 $f_{BS} \sim f_{ECCD}$ for 2X & 1O; reduced EC power for stabilization
- Confinement improvement & stationary sustainment by NTM stabilization ($\beta_N \sim 2.9$, $H_{89PL} \sim 1.8$)

Measurement of NTM structure

- Decrease/ increase in $J(r)$ during NTM growth/ suppression
- Asymmetry in δT_e during NTM stabilization
Hypothesis: 'temperature increase inside island'