



Plasma response to $m/n=3/1$ helical perturbation fields generated by the Dynamic Ergodic Divertor (DED) on TEXTOR

H R Koslowski¹, Y. Liang¹, O. Zimmermann¹, A. Krämer-Flecken¹, M. de Bock², K. H. Finken¹, R. Jaspers², E. Westerhof², R. Wolf¹, and TEXTOR-team

¹Forschungszentrum Jülich GmbH, Institut für Plasmaphysik, Association EURATOM-FZJ, D-52425 Jülich, Germany*


²FOM-Institute for Plasma Physics Rijnhuizen, Assoc. EURATOM-FOM, PO Box 1207, 3430 BE Nieuwegein, The Netherlands*

** partners in the Trilateral Euregio Cluster*


Outline

- Setup of the Dynamic Ergodic Divertor on TEXTOR
- Structure of the $m/n=3/1$ perturbation field
- Threshold for mode locking
- Onset scalings of error field induced modes
- Summary and conclusion


TEXTOR Dynamic Ergodic Divertor (DED) ..



.. consists of 16(+2) helical coils on the HFS



Schematic setup of the DED 3/1 configuration



DED parameters


- 16 coils (+2 compensation coils) mounted on the HFS
- Helical pitch resonant to $q=3$ field lines
- Configurations
 - 12/4 : PSI studies, divertor properties
 - (6/2)
 - 3/1 : Perturbation field and MHD studies
- Currents
 - up to 15kA/coil
- Frequencies
 - dc
 - low f ac (2Hz), field rotation in co- and counter current direction
 - high f ac (1 .. 10kHz)

Structure of the perturbation field


$I_p = 300\text{kA}$, $B_t = 2.25\text{T}$, $I_{\text{DED}} = 1\text{kA}$

standard plasma
equilibrium
+
vacuum DED field


$(q_a^{\text{cyl}} = 4.5)$




Amplitudes of the $n=1$ Fourier components



Pre-existing $m/n=2/1$ tearing mode locks




Typical discharge for error field experiments




- DED in 3/1 configuration
- $B_t=2.25\text{T}$, $I_p=300\text{kA}$
- $q_{a,\text{cyl}}\sim 4.5$
- co-NBI $\sim 300\text{kW}$
- at critical DED current ($\sim 0.7\text{kA}$) a 2/1 tearing mode is excited
- effect is highly reproducible
- dependence on electron density, heating power (beta), and plasma rotation has been studied

Tearing mode is created during DED phase




Electron density profile shows locked mode





Braking of toroidal rotation

toroidal rotation
decreases when
mode is created

v_{tor} profile indicates
rigid body rotation
(between $r_{q=1}$ and
 $r_{q=2}$)




Mode numbers are $m=2$ and $n=1$




mode numbers analysed with phase comparison method

Phase inversion of SXR data indicates tearing mode




Saturated island width up to 20% of minor radius



With ac DED mode is “locked” to external perturbation


TEXTOR #93993




Density scaling of the critical error field

TEXTOR #94442/43/52-56

$$B_{21} \sim I_{\text{DED}}$$




Beta dependence of mode threshold




Toroidal plasma rotation decreases mode threshold!?

TEXTOR #94422-30




tangential NBI in co-current direction
total heating power constant (P_{NBI} replaced by P_{ICRH})


Influence of plasma rotation on mode threshold



Plasma parameters are constant for rotation scan




Rotation dependence: JET results are different



B_t -scan with $q_a = \text{const}$

TEXTOR #94443/44/79/80



Summary

- TEXTOR-DED in 3/1 configuration allows to study the onset of error field generated 2/1 tearing modes
- First parameter scans were performed
- Density dependence of locked mode onset threshold agrees with other tokamaks (JET, DIII-D, C-mod)
- No clear B_t dependence of the mode threshold found
- ICRH heating (beta) has a strong stabilizing effect
- Co-NBI is found to destabilize the 2/1 mode, counter-NBI has a strong stabilizing influence
- TEXTOR data does not agree with JET results

Outlook and future work

- Continue detailed investigations of mode onset threshold vs toroidal plasma rotation
- Study B_t dependence of mode onset
- Analyze data with 1kHz ac DED, is there any influence of the rotating field?
- Understand difference to JET experiments
- TEXTOR experiments within the ITPA framework