


Fast-ion Transport by Alfvén Instabilities in DIII-D


W. Heidbrink, Y. Luo, M. Van Zeeland, M. Austin, K. Burrell, N. Gorelenkov, R. Harvey, E. Ruskov, R. White, and the DIII-D Team

Goal: Detailed measurements of fluctuations and fast ions to predict alpha transport in ITER

- Flattened fast-ion density profile
- More profile measurements
- Modeling status



Alfven Modes Degrade Fast-ion Confinement



- Volume-averaged neutron rate is below the classical TRANSP prediction during the strong Alfvén activity


- Fast-ion D_α (FIDA) diagnostic measures the spectrum of fast ions with 4 cm spatial resolution*

- FIDA "density" near ρ_{qmin} is reduced during the strong Alfvén activity

*Heidbrink, PPCF 46 (2004) 1855; Luo, RSI 78 (2007) 033505.


The Fast-ion Deficit Correlates with Alfvén Activity

- The strength of the Alfvén activity tends to increase with beam power in similar plasmas.
- The discrepancy between the classical prediction and the data is largest when the Alfvén modes are strong
- The FIDA deficit is larger than the neutron deficit



**For this comparison, the FIDA density and neutron rate are normalized by their values at 2.0 s in the 1-source shot (when Alfvén activity is undetectable).*


The Fast-ion Density Profile is Flattened



- During the strong Alfvén activity, the fast-ion density profile from FIDA is nearly flat

- The fast-ion profile inferred from the equilibrium* is also very flat

- The classical profile computed by TRANSP peaks on axis




*The kinetic EFIT equilibrium uses MSE and magnetics data to compute the pressure profile. Subtraction of the thermal pressure yields the fast-ion pressure. The FIDA data are normalized to the 1.20 s pressure profile.

My Questions (as of Fall '06)

- The FIDA and fast-ion pressure diagnostics do not measure the same thing: should their profiles be similar?
- Are the FIDA spatial profiles valid?
- Can the measured modes explain the flattened profile?


The Pressure and FIDA techniques diagnose the same parts of velocity space




- Different fast-ion diagnostics weight velocity space differently.
- The FIDA weighting shown here is for $E_\lambda = 50$ keV.
- After averaging over the distribution function $F(E,p)$ and over wavelengths, the FIDA and pressure diagnostics have similar weightings.

FIDA relative radial profile agrees well with TRANSP prediction in quiet plasmas

- Fast-ion distributions from TRANSP are dumped to the simulation code.
- Simulated profiles are higher as expected at the later time when electron density is lower.
- At the early time, FIDA profile is normalized to the simulated profile.
- At the later time, FIDA profile agrees with the simulated profile.
- Radial profile of fast-ion pressure inferred from kinetic EFITs (MSE data) are also consistent with TRANSP.



First Try at Radial Profile in Quiet Plasma showed Disagreement




- This one-source L-mode plasma has modest Alfvén activity!

- The EFIT fast-ion pressure profile differs from TRANSP as well


Perpendicular Fast-ion Acceleration at 4th Cyclotron Harmonic

- Neutron enhancement during ICH \rightarrow fast-ion acceleration
- FIDA data \rightarrow distribution function distorts
- Slight increase in bulk; perpendicular tail forms




Profiles agree well during ICH acceleration

- Fast-ion pressure exceeds classical (no RF) prediction.
- FIDA profiles show similar trends.
- The FIDA data are averaged over wavelength and time, then normalized to the no-RF profile.





Reasonable values for the spatial profile of ICH acceleration

- The profile peaks ~10 cm farther out than the nominal resonance layer.
- Calculate the expected fast-ion distribution function in CQL3D, then calculate the expected FIDA spectra and profile.
- The CQL3D prediction is close to the data.



Likely Explanation for Outshifted FIDA Signal: Orbit Effects




- Launch orbits in FIDA spatial volume with values of v_z that contribute strongly to the FIDA enhancement
- Representative orbits have turning points near the Doppler-shifted resonance layer.

Original Analysis Plan for Alfvén Mode Transport

1. Match linear NOVA-K eigenfunctions to ECE data.
2. Insert these modes (with experimental amplitudes) into ORBIT drift orbit code. Compute fast-ion transport.
3. Dump ORBIT distribution function. Use diagnostic simulation codes to predict signals. Compare with data.


The Mode Structure agrees with linear ideal MHD Theory

$n=3$ RSAE



- The MHD δT_e amplitude is scaled to match the ECE data
- Easy to match strongest TAE and RSAE modes
- More ambiguity in weaker modes. Used 11 toroidal modes.


Modes move fast ions from magnetic axis to half-radius



- Pitch-angle scattering is included @ experimental level of ~ 2 Hz
- Mode amplitudes are scaled up to investigate effect.
- The change in the distribution roughly doubles for a 3.8 ms run.


Does the "Sea" of Activity Cause Diffusive Transport?

122117, t = 410.600 ms



- ECE Data (Blue is strong)
- Many modes that constantly change

Can use *ad hoc* beam-ion diffusion in TRANSP to match experimental profile



- Used spatially uniform D_B in this initial run.
- Need a large D_B in core, smaller outside 0.6 to match experiment.

Revised Analysis Plan for Alfvén Mode Transport

1. Match linear NOVA-K eigenfunctions to ECE data. **OK**
2. Insert these modes (with experimental amplitudes) into ORBIT drift orbit code. Compute fast-ion transport **for a few milliseconds.**
- ~~3. Dump ORBIT distribution function. Use diagnostic simulation codes to predict signals. Compare with data.~~
- 3. Estimate diffusion from ORBIT run. Compare with TRANSP *ad hoc* diffusion coefficient that matches the data.**

My Questions & Answers (as of 4/07)


- The FIDA and fast-ion pressure diagnostics do not measure the same thing: should their profiles be similar? **Yes, they detect the distribution function in the same portion of velocity space.**
- Are the FIDA spatial profiles valid? **Yes, profiles analyzed the same way in quiet and ICH-heated plasmas make sense.**
- Can the measured modes explain the flattened profile? **I don't know yet.**

Backup slides

My Questions (as of 11/06)

- Are the FIDA spatial profiles valid?
- The FIDA and fast-ion pressure diagnostics do not measure the same thing: should their profiles be similar?
- Can the measured modes explain the flattened profile?

Fast-ion D_α (FIDA) Diagnostic



- A type of Charge Exchange Recombination Spectroscopy
- Use vertical view to avoid bright interferences
- Exploit large Doppler shift (measure wings of line)

- Background subtraction usually dominates uncertainty
- Achieved resolution: ~ 5 cm, ~ 10 keV, 1 ms.

