

Abstract Submitted
for the DPP01 Meeting of
The American Physical Society

Sorting Category: (Experimental)

**Comparison of Enhanced Confinement L-Mode
Regimes in JET and DIII-D With Impurity Seeding¹**

G.L. JACKSON, D.R. BAKER, T.E. EVANS, R.J. GROEBNER, D.L. HILLIS, G.R. MCKEE, M. MURAKAMI, G.M. STAEBLER, M.R. WADE, DIII-D National Fusion Facility, R. BUDNY, M. CHARLET, N.C. HAWKES, L.C. INGESSON, H.R. KOSLOWSKI, G. MADDISON, P. MONIER-GARBET, J. ONGENA, J. RAPP, F. SARTORI, J.D. STRACHAN, B. UNTERBERG, M. VON HELLERMAN, and EFDA-JET Workprogram Collaborators — Neon seeding has been used in the JET and DIII-D tokamaks to increase confinement in L-mode up to $H_{97Y} = 1.1$ producing a dramatic increase in both the ion temperature, nearly doubling in the core, and the neutron rate, even though Z_{eff} is substantially above unseeded reference discharges. In both devices, gyrokinetic analysis has shown that low k turbulence, predominantly due to ion temperature gradient (ITG) modes, was stabilized by an increase in the $E \times B$ shearing rate with impurity injection. We will discuss similarities and differences between JET and DIII-D along with gyrokinetic and transport analysis. The operational conditions for these L-mode discharges will be discussed, including reduction of MHD activity.

¹Work supported by the US DOE under contracts DE-AC03-99ER54463, DE-AC05-00OR22725, DE-AC02-76CH03073, DE-FG03-96ER54373.

- Prefer Oral Session
 Prefer Poster Session

Special instructions: JET Session

Date submitted: July 19, 2001

Electronic form version 1.4