

Abstract Submitted
for the DPP02 Meeting of
The American Physical Society

Sorting Category: 5.6.2 (Experimental)

OEDGE Analysis of DIII-D Simple-as-Possible Plasma Discharges that Approach Detachment¹

P.C. STANGEBY, J.D. ELDER, S. LISGO, U. Toronto, D.G. WHYTE, UCSD, B.D. BRAY, W.P. WEST, GA, J.G. WATKINS, SNL, M.E. FENSTERMACHER, M. GROTH, G.D. PORTER, LLNL, R.C. ISLER, ORNL, D. REITER, Juelich — A systematic study has been carried out of the simplest possible edge plasmas — no ELMs, no detachment — involving a large number of repeat, highly-diagnosed, single-null divertor, constant density, L-mode discharges on DIII-D. Three different densities were used, $\langle n_e \rangle = 2, 3, 4 \times 10^{19} \text{ m}^{-3}$. The focus of this report is on the outer target, which was particularly well diagnosed, including divertor Thomson scattering (DTS) measurements. As $\langle n_e \rangle$ was raised, the outer divertor plasma remained attached but closely approached the detached state with T_e for the DTS location closest to the target decreasing from 20 eV to 3 eV. The OEDGE (Onion-Skin Modeling + DIVIMP + EIRENE) interpretive code was employed to follow this transition, with the objective of establishing whether the controlling processes are identified and understood.

¹Supported in part by US DOE Contract DE-AC03-99ER54463, DE-FG03-95ER54294, DE-AC04-94AL85000, W-7405-ENG-48, DE-AC05-00OR22725, and NSERC Grant 250447-01.

- Prefer Oral Session
 Prefer Poster Session

P.C. Stangeby
stangeby@fusion.gat.com
University of Toronto

Special instructions: Poster 27, Edge/Divertor/Transport

Date submitted: July 19, 2002

Electronic form version 1.4