

Abstract Submitted for the Forty-Seventh Annual  
Meeting  
Division of Plasma Physics  
October 24–28, 2005, Denver, Colorado

Category Number and Subject:

Theory     Experiment

**Energetic Impurity Transport in Multispecies Gyrokinetic Plasmas,\*** C. Estrada-Mila, *UCSD*, J. Candy, R.E. Waltz, *GA* – We summarize a systematic study of the behavior of energetic impurities using the GYRO code [1]. Historically, the focus in this area has been on wave-particle interaction of energetic ions (beam ions or alpha particles) with global MHD modes (Alfvén eigenmodes or fishbones), with little or no discussion of effects arising from bulk plasma turbulence. Experimental results [2] and theoretical analyses tend to support a view where the interaction between energetic ions and a turbulent background is relatively unimportant. This is expected since at high energies particles have a large orbit width that, crudely speaking, averages over the turbulent transport coefficients. To understand this transition, we study the nonlinear gyrokinetic dynamics of (hot) helium ash as a function of the temperature ratio  $T_{\text{He}}/T_{\text{D}}$  in the range 1 to 100. Preliminary results indicate that helium confinement improves as  $T_{\text{He}}/T_{\text{D}}$  increases.

- [1] J. Candy and R.E. Waltz, *J. Comput. Phys.* **186**, 545 (2003).  
[2] S.J. Zweben, et al., *Nucl. Fusion* **40**, 91 (2000).

\*Work supported by US DOE under DE-FG03-95ER54309.

Oral     Poster