

Abstract Submitted for the Forty-Seventh Annual
Meeting
Division of Plasma Physics
October 24–28, 2005, Denver, Colorado

Category Number and Subject:

Theory Experiment

**Quantitative Studies of Nonlinear Interactions Between
Shear Flows and Turbulence in Experiment and Simulation,***

C. Holland, G.R. Tynan, *UCSD*, R.J. Fonck, G.R. McKee, *U. Wisconsin-Madison*, R.E. Waltz, J. Candy, *GA* – Using newly developed analysis algorithms, we present the first direct experimental measurements of the *nonlinear* shearing of drift-wave turbulence by a geodesic acoustic mode (GAM, a finite-frequency zonal flow). Density fluctuation measurements obtained via beam emission spectroscopy are combined with a velocity inference algorithm to provide a direct measurement of the shearing action. The measurements indicate that the shearing leads to a transfer of internal energy to smaller scales, in agreement with expectations from theory and simulation. To provide better context for these results, simulations from the GYRO gyrokinetic code are used to quantify the relative importance of zonal flow shearing vs “self-shearing” of drift-waves, and differences in turbulence shearing by “Rosenbluth-Hinton” zonal flows with zero mean frequency and finite-frequency GAMs. The implications of strong GAM shearing in the tokamak edge are also discussed.

*Work supported by US DOE under DE-FG02-04ER54758, DE-FG03-96ER54373, and DE-FG03-95ER54309.

Oral Poster