

**Abstract Submitted for the Forty-Ninth Annual Meeting
Division of Plasma Physics
November 12–16, 2007, Orlando, Florida**

Category Number and Subject:

Theory Experiment

Doppler Reflectometry Measurements of Medium Wavenumber Density Fluctuations and Zonal Flows in DIII-D,*

L. Schmitz, G. Wang, A.E. White, J. Justiniano, T.L. Rhodes, W.A. Peebles, *UCLA* – Doppler reflectometry is a versatile diagnostic for poloidal plasma flow measurements and local density fluctuation spectra. Depending on the launch angle and frequency of the probing beam, the signal back-scattered from the plasma cut-off layer is sensitive to density fluctuations at a specific poloidal wavenumber k_θ ($3 \text{ cm}^{-1} < k_\theta < 9 \text{ cm}^{-1}$, calculated using GENRAY ray tracing code). The plasma flow velocity v_θ is obtained with high time resolution from the Doppler shift ω_D of the back-scattered signal ($v_\theta = \omega_D/k_\theta$). Doppler reflectometry is well-suited for the detection of zonal flows, characterized by poloidal flow fluctuations ($v_\theta = k_r \Phi/B_\phi$). Zonal flows are thought to regulate the local turbulence level and radial correlation. We present first reflectometry measurements of geodesic acoustic modes (GAMs) and low frequency zonal flows in DIII-D L-mode plasmas ($0.6 < r/a < 0.9$). The interaction of these time-dependent plasma flows with medium wavenumber density fluctuations is investigated in order to study turbulence self-organization.

*Supported by US DOE under DE-FG03-01ER54615 and DE-FC02-04ER54698.