

**Abstract Submitted for the 50th Annual Meeting  
Division of Plasma Physics  
November 17–21, 2008, Dallas, Texas**

Category Number and Subject:

Theory     Experiment

**Simulation of DIII-D Plasma Shutdown by Deuterium Dilution Cooling\*** V.A. Izzo, *UCSD*, P.B. Parks, W. Wu, *GA* — To mitigate ITER disruptions and avoid large numbers of runaway electrons, a significant increase in the total (free + bound) electron inventory is likely required. The Rosenbluth criterion— $E_c \approx 0.12n_{e,20}$ —determines the critical electric field (in V/m) at which exponential runaway avalanching will occur. Here we consider instantaneous dilution cooling of a DIII-D plasma by the injection of 100 times the initial deuterium density to simulate rapid core penetration of a  $D_2$  pellet train or liquid jet. The 3D NIMROD MHD simulation is initialized with an equilibrium pressure profile, but a  $100 \times$  density increase and a corresponding  $100 \times$  temperature reduction. The plasma is assumed to have in situ carbon fraction of 1% of the pre-dilution density, which produces strong edge radiation at the dilution cooled temperatures. A cooling front propagates inward and ultimately triggers a central 1/1 MHD event. The central current density transiently increases by more than a factor of 2. The 3D simulation is compared with a 1D simulation which incorporates a Kadomtsev mixing model for the 1/1 MHD event.

\*Work supported by US DOE under DE-FG03-95ER54309.