

**Abstract Submitted for the 50th Annual Meeting  
Division of Plasma Physics  
November 17–21, 2008, Dallas, Texas**

Category Number and Subject:

Theory     Experiment

**Long-Wavelength Turbulence Scaling Properties in DIII-D,\***  
G.R. McKee, D.J. Schlossberg, M.W. Shafer (UW-Madison), C.H. Holland (UCSD), P. Gohil (GA) – The scaling properties of long-wavelength density fluctuations are investigated in DIII-D L-mode and H-mode plasmas utilizing the expanded high-sensitivity 2D Beam Emission Spectroscopy (BES) system. BES employs a 64-channel system that utilizes a radially-scannable 8x8 array sampling multiple radial and poloidal correlation lengths, allowing for full sampling of the 2D wavenumber spectrum. Measurements of turbulence as a function of several important dimensionless parameters ( $\kappa$ ,  $T_e/T_i$ , ion mass,  $\rho_*$ ) are obtained, showing that fluctuation intensity increases strongly with decreasing plasma elongation (at constant  $q$ ), consistent with increased thermal transport and reduced energy confinement. In contrast, increasing  $T_e/T_i$  increases momentum and thermal transport with little change in low- $k$  density fluctuations. Measurements obtained during a  $\rho_*$  ( $\rho_i/a$ ) scan in hydrogen will also be presented. Together, these measurements will be crucial for comparing with transport simulations, such as GYRO and TGLF.

\*Work supported by the US DOE under DE-FG02-89ER53296, DE-FG02-97ER54917, and DE-FC02-04ER54698.