

**Abstract Submitted for the 50th Annual Meeting
Division of Plasma Physics
November 17–21, 2008, Dallas, Texas**

Category Number and Subject: 5.6.2. DIII-D Tokamak

☐ Theory ☒ Experiment

Phase Contrast Imaging Measurements of Turbulence in ELMing Plasmas,* J.C. Rost, M. Porkolab, J.R. Dorris, *Plasma Science and Fusion Center, MIT*, K.H. Burrell, *General Atomics* – The Phase Contrast Imaging (PCI) diagnostic on DIII-D has been upgraded since the last study of ELMs. Signal-to-noise and bandwidth have been increased, and a rotating mask system added to spatially localize measurements of electrostatic modes is used here to determine if modes are aligned with the magnetic field. During Type I ELMs, a burst of turbulence coincides with the magnetic perturbation but lasts longer, several milliseconds compared to a few 100 microseconds. ELM-like modes seen between Type I ELMs (possibly Type II ELMs) are smaller and shorter on the magnetics and D_{α} , but the density perturbations on the PCI are similar in amplitude to that during the Type I ELMs. The turbulence during the ELM-like modes has a phase velocity about one third that of the turbulence during the Type I ELM, indicating a different radial location or different driving instability. Measurements with the mask show that the ELM-related turbulence is aligned with the total background magnetic field, similar to electrostatic modes.

*Work supported in part by the US DOE under DE-FC02-99ER54512 and DE-FC02-04ER54698.