

**Abstract Submitted for the 51st Annual Meeting
Division of Plasma Physics
November 2–6, 2009, Atlanta, Georgia**

Density Fluctuation Enhancement During RMP ELM-Suppressed Discharges,* G.R. McKee, R.J. Fonck, D.J. Schlossberg, M.W. Shafer, Z. Yan, *U. Wisc.*; T.E. Evans, *GA*; R.A. Moyer, *UCSD* — Fluctuation behavior changes dramatically over the radial range $0.6 < r/a < 1.0$ during ELM-suppressed phases of H-mode discharges with externally applied resonant magnetic perturbations (RMPs) on DIII-D. The 2D spatiotemporal characteristics of density fluctuations during ELM suppression are characterized with beam emission spectroscopy and exhibit a broadband structure from 80-400 kHz. The fluctuation power increases by up to one order of magnitude in the mid-core region ($0.6 < r/a < 0.8$), with more modest increases in the pedestal region. The increase in turbulence coincides with the ELM-suppressed phases, which typically start a few 100 ms after the application of a radial magnetic field. The radial and poloidal correlation lengths of the increased fluctuations are a few cm., typical of long-wavelength turbulence. The increased fluctuations appear to be related to observed profile changes, reduction of the line integrated density, and increased radial particle transport.

*Work supported by the US DOE under DE-FG02-89ER53296, DE-FG02-08ER54999, DE-FC02-04ER54698 and DE-FG02-07ER54917.