

**Abstract Submitted for the 51st Annual Meeting
Division of Plasma Physics
November 2–6, 2009, Atlanta, Georgia**

Measurements of the Wavenumber Spectrum of Multi-scale Turbulence in the Core of DIII-D H-mode Plasmas* L. Schmitz, G. Wang, J.C. Hillesheim, W.A. Peebles, T.L. Rhodes, E.D. Doyle, L. Zeng, *UCLA*, C. Holland, *UCSD*, A.E. White, *ORISE*, G.R. McKee, *U. Wisc.*, J.C. DeBoo, J.C. deGrassie, K.H. Burrell, C.C. Petty, *GA* — The wavenumber spectrum and spectral index of multi-scale turbulence ($0.5 \leq k_{\theta} \rho_s \leq 6$) in the core of L and H-mode DIII-D plasmas have been determined by Doppler Backscattering (DBS). A comparison is made to quasilinear spectra from the trapped gyro-Landau fluid code (TGLF) to assess the contributions of ITG and TEM/ETG turbulence to transport fluxes ($0.4 \leq r/a \leq 0.8$). Initial results of comparisons to spectra calculated via nonlinear gyrokinetic (GYRO) simulations are also presented. In high temperature, low density plasmas, ITG-scale and intermediate-scale core turbulence is found reduced by at least an order of magnitude across the L- to H-mode transition ($T_i/T_e \geq 2$). This reduction is attributed to the combined effects of reduced turbulence drive and increased core ExB flow shear in H-mode.

*Work supported by the US DOE under DE-FG03-08ER54984, DE-FG02-07ER54917, DE-AC05-06OR23100, DE-FG02-89ER53296, and DE-FC02-04ER54698.