

**Abstract Submitted for the 52nd Annual Meeting
Division of Plasma Physics
November 8–12, 2010, Chicago, Illinois**

Category Number and Subject: 5.1.0 Turbulence & Transport

Theory Experiment

Characterization of Neoclassical Transport in the Plasma Edge,* E.A. Belli, J. Candy, *General Atomics* – An extensive characterization of neoclassical transport in the plasma edge is studied numerically using the NEO code, which solves a hierarchy of drift-kinetic equations based on an expansion in ρ_{*i} . The code includes the self-consistent coupling of electrons and multiple ion species, fully general geometry, including up-down asymmetry effects, and full rapid toroidal rotation effects. The validity of standard local neoclassical transport theory in the H-mode edge is assessed for typical DIII-D plasmas via solution of the higher-order drift-kinetic equations, and the influence of finite-orbit-width effects on the non-local neoclassical energy transport and flows is explored. Preliminary results indicate that only a weak finite-orbit-width effect is found due to the steep gradients and thus the δf formulation is valid for most of the pedestal. Extended studies of the influence of impurities and orbit-squeezing effects on the edge transport are presented. Limitations of analytic theories, such as the Chang-Hinton theory and the Sauter model, are also studied for realistic experimental parameters.

*Supported by US DOE under DE-FG02-95ER54309 and DE-FC02-06ER54873