

**Abstract Submitted for the 54th Annual Meeting
Division of Plasma Physics
October 29 through November 2, 2012
Providence, Rhode Island**

Category Number and Subject: 5.6.2. DIII-D Tokamak &
5.7.0 Diagnostics

[] Theory [**X**] Experiment

Bursts in Electron Cyclotron Emission During ELMs and EHOs in Tokamaks*, E. Li, M.E. Austin, *University of Texas-Austin*
- Electron cyclotron emission (ECE) bursts during edge localized modes (ELMs) have been observed and reported in several tokamaks. A number of possible explanations have been suggested, but none describe completely the observed ECE bursts. A key to understanding these bursts is that they always occur during edge MHD activity such as ELMs or EHOs in QH-mode plasmas. Other observed properties are that the ECE bursts appear on the low-field-side with an intensity of several tens or hundreds times the thermal emission level, and with a bandwidth as narrow as the instrumental filter spacing of 400 MHz or narrower. A model based on the basic emission and absorption relationship of ECE in plasma is put forth. We calculate the radiation temperature for different electron velocity distribution functions, but with a small variable resonance frequency within the instrumental bandwidth. The resulting radiation temperature and its spectral width agree well with experimental observations both in non-Maxwellian and Maxwellian distribution, but the radiation temperature in non-Maxwellian case is much higher.

* This work was supported in part by the US Department of Energy under DE-FG03-97ER54415.