

**Abstract Submitted for the 54th Annual Meeting
Division of Plasma Physics
October 29 through November 2, 2012
Providence, Rhode Island**

Category Number and Subject: 5.6.2. DIII-D Tokamak

[] Theory [X] Experiment

Measurements and Modeling of the Divertor Heat Flux Width in DIII-D,* M.A. Makowski, C.J. Lasnier, *LLNL*; J. Nichols, *PPPL*; A.W. Leonard, T.H. Osborne, P.B. Snyder, *General Atomics* – The relationship between measured divertor heat flux and upstream SOL profiles is explored to test critical pressure gradient models of radial energy transport in DIII-D. We are developing a physics based model that extends the well validated kinetic ballooning mode pedestal paradigm up to the separatrix and into the SOL. Evidence in support of this model is that the measured separatrix pressure gradient scales similarly to the critical pressure gradient for the infinite- n ballooning mode obtained from the BALOO code. Recent results have revealed how the heat flux width scales with plasma parameters [1]. The main result is that $\lambda_{sol} \sim B_p^{-1}$ and is independent of all other parameters including machine size. Here λ_{sol} is the heat flux width in the scrape-off-layer (SOL) and B_p is the poloidal field at the outer mid-plane. New measurements of the heat flux width versus the injected power demonstrate a weak dependence, $\lambda_{sol} \sim P_{inj}^{-0.1}$, consistent with other experiments.

[1] T. Eich et al., Phys. Rev. Lett. **107**, 215001 (2011); M.A. Makowski, Phys. Plasmas **19**, 056122 (2012).

*This work performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under DE-AC52-07NA27344 and in part by the U.S. Department of Energy under DE-AC02-09CH11466 and DE-FC02-04ER54698.