

Abstract Submitted for the 56th Annual Meeting
Division of Plasma Physics
October 27–31, 2014
New Orleans, Louisiana

Category Number and Subject: DIII-D Session

Theory Experiment

RMP-Induced Plasma Transport Near X point,* J.D. Callen, C.C. Hegna, *U. Wisconsin-Madison* – The experimentally most noticeable effect of resonant magnetic perturbations (RMPs) on H-mode plasmas is "density pump-out." This effect is most evident in the near-separatrix region where RMPs can cause the electron density gradient there to decrease by a factor of up to two. The previously developed flutter model [1] produces some RMP-induced transport in this region. However, two other electron effects need to be taken into account in the X point region of the divertor separatrix. First, small 3D fields can cause significant (many cm) radial motion of field lines in "homoclinic tangles" very near the X point which have been observed experimentally in DIII-D. A flutter-type plasma transport model based on parallel electron collisional effects caused by RMP-induced "radial" motion of field lines away from the lowest order axisymmetric magnetic flux surfaces in the X point region is being developed. The second effect is that a small fraction of long length magnetic field lines in the near-separatrix region are "open" ones which are directly connected to material walls in the divertor region. Electrons on such field lines could conduct significant electron heat to the divertor plates.

[1] J.D. Callen, et al., *Nucl. Fusion* **53**, 113015 (2013).

*Work supported by the US DOE under DE-FG02-92ER4139 and DE-FG02-86ER53218.