

Abstract Submitted for the 56th Annual Meeting
Division of Plasma Physics
October 27–31, 2014
New Orleans, Louisiana

Category Number and Subject:

[] Theory [X] Experiment

3D Field-Induced Transport and Plasma Response Leading to ELM Suppression in DIII-D,* S.P. Smith, C. Paz-Soldan, R.J. Groebner, *GA*; O. Meneghini, *ORAU*; R. Nazikian, B.A. Grierson, *PPPL*; M.E. Austin, *U Texas*; J.D. Callen, *U Wisc.-Madison*; E.M. Davis, *MIT*; R.A. Moyer, *UCSD*; T.L. Rhodes, G. Wang, L. Zeng, *UCLA* – A clear increase in trapped electron mode (TEM) scale density fluctuation levels \tilde{n}_e is seen at the top of the pedestal as the plasma transitions from edge localized mode (ELM)ing to ELM suppression with applied 3D resonant fields. Additional increases in T_e fluctuations and libe-integrated n_e at the top of the pedestal are seen as the 3D field strength is increased. High resolution T_e and n_e profile measurements near the top of the pedestal show strong transport scaling with the applied field ($1/L_{Te}, 1/L_{ne} \sim I_c^2$) during ELM suppression. These latter results are consistent with the magnetic flutter model regulating transport at the top of the pedestal, possibly driven by kink mode coupling, however the former results support a 3D modification of microturbulence stability as the process by which ELMs are suppressed.

*Work supported by the US Department of Energy under DE-FC02-04ER54698, DE-AC02-09CH11466, DE-FG03-97ER54415, DE-FG02-92ER54139, DE-FG02-94ER54235, DE-FG02-07ER44917, and DE-FG02-08ER54984.