

**Abstract Submitted for the 56th Annual Meeting  
Division of Plasma Physics**

Category Number and Subject: 6.20 DIII-D Tokamak

Theory     Experiment

**Using Quiescent H-mode to Access an Improved High Pressure Plasma Edge,\*** W.M. Solomon, B.A. Grierson, R. Nazikian, *PPPL*; K.H. Burrell, A.M. Garofalo, T.H. Osborne, P.B. Snyder, *GA*; A. Loarte, *ITER*; G.R. McKee, *U Wisc*; M.E. Fenstermacher, *LLNL* – Experiments on DIII-D have extended Quiescent H-mode (QH-mode) to high density through the use of strong shaping, overcoming a long-standing limitation in QH-mode operation, a high confinement state of the plasma that does not exhibit edge localized modes. These experiments have navigated a valley of improved edge peeling-ballooning stability dubbed “Super H-mode”, which opens up at high density with strong plasma shaping. The thermal energy confinement time increases due to improvements in both the pedestal height and the core transport. Theoretical calculations of the pedestal height and width as a function of density using the EPED model are in quantitative agreement with the measurements. Together with the achievement of high beta, high confinement and low  $q_{95}$  for many energy confinement times, these results extend QH-mode as a potentially attractive operating scenario for ITER and point to a path for a new high performance regime that could improve the attractiveness of a fusion reactor.

\*Work supported by the US DOE DE-AC02-09CH11466, DE-FC02-04ER54698, DE-FG02-89ER53296, DE-FG02-08ER54999 and DE-AC52-07NA27344.