

Abstract Submitted for the 56th Annual Meeting
Division of Plasma Physics
October 27–31, 2014
New Orleans, Louisiana

Category Number and Subject:

Theory Experiment

Development and Validation of a Critical Gradient Energetic Particle Driven Alfvén Eigenmode Transport Model for DIII-D Tilted Neutral Beam Experiments,* R.E. Waltz, GA;

and E.M. Bass, *UCSD* —Recent experiments on DIII-D with tilted neutral beam injection (NBI), which significantly vary the beam energetic particle (EP) source profiles, have provided strong evidence that unstable Alfvén eigenmodes (AE) drive stiff EP transport at a critical EP density gradient [1]. We hope to identify the critical gradient with the condition that the maximum local AE growth rate falls to the local ion temperature gradient (ITG)/trapped electron mode (TEM) rate at the same low toroidal mode number. This condition was supported by early nonlinear local GYRO simulations [2] and more is more optimistic than stiff EP transport at the AE marginal stability gradient used in a recent ITER projection of AE driven alpha confinement losses [3]. The AE and ITG/TEM growth rates are from GYRO with comparison of Maxwellian and slowing down beam-like EP distributions.

[1] W.W. Heidbrink, *et al.*, *Nucl. Fusion* **53**, 093006 (2013).

[2] E.M. Bass and R.E. Waltz, *Phys. Plasmas* **17**, 112319 (2010).

[3] E.M. Bass, *Bull. Am. Phys. Soc.* **58**, 168 (2013).

*Work supported by the US DOE under DE-FG02-95ER54309 and DE-FC02-08ER54977.