

**Abstract Submitted for the 56th Annual Meeting
Division of Plasma Physics
October 27–31, 2014
New Orleans, Louisiana**

Category Number and Subject:

Theory Experiment

TGLF Analysis of ITG-TEM Transitions in DIII-D Plasmas,* X. Wang, S. Mordijck, *William & Mary*; G.M. Staebler, *GA*; O. Meneghini, *ORAU* – TGLF [1] calculations show that, when adding electron cyclotron heating, DIII-D H-mode plasmas can transit from the ion temperature gradient (ITG) to trapped electron mode (TEM) domain around mid-radius. With different turbulence scale-lengths as well as opposite drift direction, ITG/TEM plasmas have different transport features, and thus lead to different density and rotation profiles. Detailed TGLF sensitivity analysis shows that the trapped electron mode's growth-rate is more sensitive to electron temperature gradient than the density gradient. Furthermore, by varying the input torque from co-IP to counter-IP through neutral beams, the ITG/TEM reversing radial location changes. We compare three parameters: density fluctuations, linear instability growth-rates, and ExB shearing rates in these discharges. We show that, for co- and counter-beam discharges, the ExB shearing rate becomes comparable or even larger than the growth-rate in the plasma edge. However, under balanced beam injection, the growth-rate is always larger than the shearing rate..

[1] G.M. Staebler, et al. *Phys. Plasmas* **12**, 102508 (2005).

*Work supported by the US Department of Energy under DE-SC0007880 and DE-FG02-95ER54309.