

Abstract Submitted
for the DPP98 Meeting of
The American Physical Society

Sorting Category: 5.1.1.2 (experimental)

Power Dependence of the 2D Divertor Emission Profiles During the Transition to PDD Operation in DIII-D¹

M.E. FENSTERMACHER, S.L. ALLEN, C.J. LASNIER, R.D. WOOD, Lawrence Livermore National Laboratory, A.W. LEONARD, T.W. PETRIE, General Atomics, R.J. COLCHIN, R.C. ISLER, Oak Ridge National Laboratory — The time histories of 2D spatial profiles of radiated power (P_{rad}) and the visible emission from carbon and deuterium are presented during the transition from attached ELMing H-mode to Partially Detached Divertor (PDD) operation. Previous work² showed that in high power (12 MW) rapidly ELMing (250 Hz) discharges the D_{α} emission at the beginning of the transition increased first in the private flux region (PFR) near the inner strike point and then filled the PFR; carbon increased first along the outer leg separatrix and then near the X-point. This paper compares transitions at high and low power (ELM frequency). Animations of the 2D profiles for P_{rad} , the constituent radiators, and the regions dominated by deuterium recombination will be shown. The 2D spatial profile of carbon radiated power will be determined by normalizing the visible CIII profiles with absolutely calibrated chordal measurements of VUV lines in the divertor.

¹Work supported by U.S. DOE Contracts W-7405-ENG-48, DE-AC03-89ER51114, and DE-AC05-96OR22464.

²M.E. Fenstermacher *et al.*, to be published in J. Nucl. Mater. (1999).

- Prefer Oral Session
 Prefer Poster Session

Max E. Fenstermacher
fenstermache@gav.gat.com
Lawrence Livermore National Laboratory

Special instructions: DIII-D Oral Session II, immediately following Wade

Date submitted: July 20, 1998

Electronic form version 1.3