

Abstract Submitted
for the DPP98 Meeting of
The American Physical Society

Sorting Category: 5.7 (experimental)

**Studies of Bubble Neutron Detectors For Alpha
Knock-on Tail Measurements¹**

R.K. FISHER, P.B. PARKS, General Atomics, S.S. MEDLEY, A.L. ROQUEMORE, Princeton Plasma Physics Laboratory, M.L. LOUGHLIN, JET Joint Undertaking, E. MORSE, A. BELIAN, University of California, Berkeley, J. LIPTAC, University of Washington — Measurement of the DT neutron energy spectrum above 16 MeV will yield information on confined alphas in ITER. These neutrons result from fusion reactions involving energetic ions created by knock-on alpha-fuel ion collisions. Bubble neutron detectors, designed to only detect neutrons with energies above a selectable threshold energy, were used for preliminary measurements of the high energy neutron spectra from TFTR and JET DT plasmas. Since there was not time to calibrate the detectors prior to the DT tokamak experiments, recent work has concentrated on measuring the detector response as a function of neutron energy at accelerator neutron sources at Ohio University and at UC Berkeley. Results from the accelerator tests will be reported and used to interpret the tokamak results, and to plan for alpha diagnostics on future experiments including JET DTE2 and ITER.

¹Work supported by U.S. DOE under Grant DE-FG03-92ER54150 and Contract DE-AC02-76CH03073.

Prefer Oral Session
 Prefer Poster Session

R.K. Fisher
fisherr@gav.gat.com
General Atomics

Date submitted: July 21, 1998

Electronic form version 1.3