

Abstract Submitted
for the DPP98 Meeting of
The American Physical Society

Sorting Category: 5.1.1.2 (experimental)

**The Dependence of Local Edge Plasma Parameters
on Heating Power at Partial Detachment Onset in DIII-D¹**

R. MAINGI, M.R. WADE, Oak Ridge National Laboratory, M.E. FENSTERMACHER, Lawrence Livermore National Laboratory, A.W. LEONARD, M.A. MAHDAVI, T.W. PETRIE, General Atomics, J.G. WATKINS, Sandia National Laboratories — We reported² a study of plasma density limiting processes and successful achievement of discharges with density $\sim 50\%$ above the empirical Greenwald density limit with H-mode confinement. Because total detachment is the first and most common density limit process in DIII-D, we have measured the local upstream separatrix electron density and temperature ($n_e^{\text{sep,det}}$, $T_e^{\text{sep,det}}$) at detachment onset (partial detachment) as a function of heating power, P_{heat} to compare with predictive models. We find that $n_e^{\text{sep,det}} \propto P_{\text{heat}}^{0.8}$ and $T_e^{\text{sep,det}} \propto P_{\text{heat}}^{0.6}$. Our data are in reasonable agreement with Borass' equations³ used to model partial detachment onset, which predict that the $n_e^{\text{sep,det}} \propto P_{\text{heat}}^{0.6-0.7}$ and $T_e^{\text{sep,det}} \propto P_{\text{heat}}^{0.3}$.

¹Work supported by U.S. DOE under Contracts DE-AC05-96OR22464, W-7405-ENG-48, DE-AC093-89ER51114, and DE-AC04-94AL85000.

²R. Maingi *et al.*, Phys. Plasmas **4** (1997) 1752.

³K. Borass, Nucl. Fusion **31** (1991) 1035.

Prefer Oral Session
 Prefer Poster Session

R. Maingi
maingi@gav.gat.com
Oak Ridge National Laboratory

Special instructions: DIII-D Oral Session II, immediately following Colchin

Date submitted: July 21, 1998

Electronic form version 1.3